

First karyotype description of the species of *Adenomera* Steindachner, 1867 (Anura, Leptodactylidae) in the “*thomei*” clade

Ramon Costa Dominato^{1,2}, Guilherme Costa de Oliveira^{2,4},
Carla Santana Cassini², Victor Goyannes Dill Orrico^{2,3},
Cléa dos Santos Ferreira Mariano^{3,4,5}, Janisete Gomes Silva^{1,3,4}

1 Programa de Pós-Graduação em Genética e Biologia Molecular, Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz, Rodovia Jorge Amado, Km 16. CEP: 45662-900 Ilhéus, BA Brazil
2 Tropical Herpetology Laboratory, Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz, Rodovia Jorge Amado, Km 16. CEP: 45662-900 Ilhéus, BA Brazil
3 Programa de Pós-Graduação em Zoologia, Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas, Rodovia Jorge Amado, Km 16. CEP 45662-900 Ilhéus, BA Brazil
4 Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz, Rodovia Jorge Amado, Km 16. CEP 45662-900 Ilhéus, BA Brazil
5 Laboratório de Artrópodes Sociais - LABAS Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz, Rodovia Jorge Amado, Km 16. CEP: 45662-900 Ilhéus, BA Brazil

Corresponding author: Ramon Costa Dominato (ramondominato@gmail.com)

Academic editor: Syed Farhan Ahmad | Received 23 February 2022 | Accepted 6 April 2022 | Published 30 August 2022

<http://zoobank.org/1A863C5A-777C-4A14-9A72-2A6E32D6AAEA>

Citation: Dominato RC, de Oliveira GC, Cassini CS, Orrico VGD, Mariano CSF, Silva JG (2022) First karyotype description of the species of *Adenomera* Steindachner, 1867 (Anura, Leptodactylidae) in the “*thomei*” clade. *Comparative Cytogenetics* 16(3): 151–159. <https://doi.org/10.3897/compcytogen.v16.i3.82641>

Abstract

The genus *Adenomera* Steindachner, 1867 currently comprises 29 nominal species, some of which are suggested to be cryptic species complexes. The present study was carried out with specimens of the “*thomei*” clade that encompasses three taxa distributed in the Atlantic Forest biome: *Adenomera thomei* Almeida et Angulo, 2006, *Adenomera* sp. L., and *Adenomera* sp. M. We used classical cytogenetics to describe the diploid number and karyomorphology of these three species collected in two different locations in the state of Bahia, Brazil. Our results revealed the diploid number $2n = 24$ (FN = 34) with two pairs of metacentric chromosomes (pairs 1 and 5), three pairs of submetacentric chromosomes (pairs 2, 3, and 4), and seven pairs of telocentric chromosomes (pairs 6, 7, 8, 9, 10, 11, and 12). Further morphological, bio-acoustic, and cytogenetic data (C-banding and AgNor) are needed to better delineate the lineages within the “*thomei*” clade.

Keywords

Chromosomes, cryptic species, cytogenetics, Giemsa, taxonomy

Introduction

The genus *Adenomera* Steindachner, 1867 currently comprises 29 described species that are distributed from tropical South America to the east of the Andean region (Carvalho et al. 2021). Due to the history of systematic reviews and the complex taxonomy of this group, taxonomic knowledge has not kept pace with the knowledge on its phylogeny (Duellman 2005; Menin et al. 2008; Fouquet et al. 2014). Out of the several hurdles for taxonomic studies on this genus, we highlight the high intra and interspecific similarities and the presence of cryptic species complexes (Fouquet et al. 2014). The difficulty increases when studies use only molecular data, disregarding other characteristics and making the interpretation of results less accurate (Pyron and Wiens 2011; De Sá et al. 2014).

Cytogenetic studies on the genus *Adenomera* date from the 1970s (Bogart 1970, 1974) when the karyotypes of *Adenomera andreae* (Müller, 1923), *Adenomera hylaedactyla* (Cope, 1868), *Adenomera lutzi* Heyer, 1975, and *Adenomera marmorata* (Steindachner, 1867) were described. However, the volume of cytogenetic information for the genus has not significantly advanced over these five decades. Campos et al. (2009) described the karyotypes of individuals from western São Paulo associating them with the nominal species *Adenomera* aff. *bokermanni* Heyer, 1973, *A. hylaedactyla*, and *A. marmorata*. Additionally, the karyotype of *Adenomera diptyx* (Boettger, 1885) was described by Zaracho and Hernando (2011). Thus, there is cytogenetic information for only five species among the 29 species described for this genus. Therefore, the small number of described karyotypes makes it difficult to both understand the chromosomal evolution of the genus and to better delimit species (Campos et al. 2009; Zaracho and Hernando 2011).

Among the clades within the genus *Adenomera*, the species of the *thomei* clade, *Adenomera thomei* Almeida and Angulo (2006), *Adenomera* sp. L, and *Adenomera* sp. M, are restricted to the Atlantic Forest in Brazil. *Adenomera thomei* was described from specimens collected in a cocoa plantation in the municipality of Linhares in the state of Espírito Santo (Almeida and Angulo 2006). Currently, there are records of this species also in the states of Rio de Janeiro, São Paulo, Minas Gerais, and Bahia (Almeida and Angulo 2006; Fouquet et al. 2014). The specific boundaries among these lineages are unclear mainly due to the lack of information on *Adenomera* sp. L and *Adenomera* sp. M, both of which are found only in the southern region of the state of Bahia (Fouquet et al. 2014). Knowledge on the bioacoustics, morphology, and cytogenetics for representatives of this clade is scarce (Angulo et al. 2003; Angulo 2004; Duellman 2005) and thus far it has not been used to distinguish between these two lineages.

Karyotypic information associated to DNA sequence data has helped clarify the taxonomy and systematics of some Brazilian anuran groups (Lourenço et al. 2008; Targueta et al. 2010; Suárez et al. 2013; Lourenço et al. 2015; Ferro et al. 2018; Marciano-Jr et al. 2021). To date, all information available regarding cytogenetic data within *Adenomera*

populations is taxonomically inconclusive (e.g., Campos 2009). Nevertheless, these chromosome data provided support on taxonomic decisions on a broad study of species delimitation of *Adenomera marmorata*, which included DNA sequence, morphological, and bioacoustic data (Cassini et al. 2020). Thus, it is clear that further cytogenetic studies on the genus *Adenomera* will allow more robust conclusions regarding this taxonomically challenging group. The objective of this study was to describe for the first time the karyotype of *Adenomera* species of the “*thomei*” clade from different locations in southern Bahia and compare the chromosomal patterns among the specimens.

Material and methods

Cytogenetic analysis was performed using 12 specimens of two species in the “*thomei*” clade collected in three sites in the state of Bahia (BA) (Table 1) under the SISBIO license 62181. The specimens were taken to the Tropical Herpetology Laboratory at the Universidade Estadual de Santa Cruz (UESC), Ilhéus, Bahia, Brazil. We identified the specimens collected in the municipality of Ilhéus as *Adenomera* cf. *thomei*, since the bioacoustic data showed the same pattern as that recorded for populations in the “*thomei*” clade.

Table 1. Information on *Adenomera* specimens in the “*thomei*” clade used in this study.

Voucher	Genus	Species	Sex	Locality	Coordinates
MZUESC 22146	<i>Adenomera</i>	cf. <i>thomei</i>	Juvenile	Ilhéus - BA	-14.800189, -39.154594
MZUESC 22147	<i>Adenomera</i>	cf. <i>thomei</i>	Juvenile	Ilhéus - BA	-14.800189, -39.154594
MZUESC 22148	<i>Adenomera</i>	cf. <i>thomei</i>	Juvenile	Ilhéus - BA	-14.795269, -39.037339
MZUESC 22149	<i>Adenomera</i>	cf. <i>thomei</i>	Male	Ilhéus - BA	-14.795269, -39.037339
MZUESC 22150	<i>Adenomera</i>	cf. <i>thomei</i>	Male	Ilhéus - BA	-14.795269, -39.037339
MZUESC 22151	<i>Adenomera</i>	sp. L	Male	Igrapiúna - BA	-13.821933, -39.171175
MZUESC 22152	<i>Adenomera</i>	sp. L	Juvenile	Igrapiúna - BA	-13.821933, -39.171175
MZUESC 22153	<i>Adenomera</i>	sp. L	Male	Igrapiúna - BA	-13.821933, -39.171175
MZUESC 22154	<i>Adenomera</i>	sp. L	Female	Igrapiúna - BA	-13.821933, -39.171175
MZUESC 22155	<i>Adenomera</i>	sp. L	-	Igrapiúna - BA	-13.821933, -39.171175
MZUESC 22156	<i>Adenomera</i>	sp. L	-	Igrapiúna - BA	-13.821933, -39.171175
MZUESC 22157	<i>Adenomera</i>	sp. L	Juvenile	Igrapiúna - BA	-13.821933, -39.171175

We followed the protocol of Schmid (1978) with modifications. In the present study, a 2% colchicine solution (0.1 ml/10 g of weight) was used during 4–6 h. Subsequently, the specimens were sacrificed with lidocaine gel at a concentration of 5% spread over the entire body. The vouchers were fixed in 10% formaldehyde for 24 hours, kept in 70% alcohol, and deposited at the UESC Herpetological collection.

Chromosomal preparations were obtained from intestinal cells. The intestinal epithelium was kept in a hypotonic solution (0.075 M KCL) for 40 minutes and fixed in CARNOY solution (3:1 methanol: acetic acid). Then, the cell suspensions were placed on the surface of a slide and dried at room temperature in the dark. To determine chromosome composition and the fundamental number (FN), cells were stained with 3% Giemsa for 10 minutes. Chromosomes were classified according to Green and Sessions (1991) as metacentric (M), submetacentric (SM), subtelocentric (ST), and telocentric (T) (Table 2).

Results obtained were compared with cytogenetic data available in the literature. The images were captured and analyzed using an Olympus BX-51 microscope, a Q-Capture Pro image capture camera, and the Image Pro Plus software. We used Adobe Photoshop CC 2019 for the analysis and arrangement of the karyotype in descending order.

Table 2. *Adenomera* species with described karyotype, fundamental number and bibliographic references. Species Identification followed the taxon name used in the original contribution.

Species	Karyomorphology	Diploid number	Fundamental number	References
<i>A. diptyx</i>	1M+3SM+ 9T	26	FN = 34	Zaracho and Hernando 2011
<i>A. andreae</i>	1M+4SM+2ST, 6T	26	FN = 40	Bogart 1974
<i>A. lutzi</i>	-	26	NA	Bogart 1970 apud Kuramoto, 1990
<i>A. hylaedactyla</i>	1M+ 3SM+ 9T	26	FN = 34	Campos et al. 2009
<i>A. hylaedactyla</i>	1M+1SM+2ST+9T	26	FN = 36	Bogart 1974
<i>A. marmorata</i>	2M+1SM+2ST+7T	24	FN = 34	Bogart 1974
<i>A. cf. marmorata</i>	3M+3SM+6T	24	FN = 34	Campos et al. 2009
<i>A.cf. marmorata</i>	2M+3SM+7T	24	FN = 34	Campos et al. 2009
<i>Adenomera</i> sp. L	2M+3SM+7T	24	FN = 34	Present Study
<i>A. cf. thomei</i>	2M+3SM+7T	24	FN = 34	Present Study
<i>A. cf. bokermanni</i>	2M+3SM+1ST+4T+3NP (1M + 2T)	23	FN = 34	Campos et al. 2009

Results

We analyzed metaphases of 12 individuals of the lineages *Adenomera* sp. L (n = 6) and *Adenomera* cf. *thomei* (n = 6; sex of specimens is shown in Table 1). The karyotype of all analyzed specimens showed 2n = 24 (FN = 34) and no heteromorphic sex chromosomes. All individuals showed the karyotype 2n = 24 with a karyotypic formula of 4M + 6SM + 14T (metacentric pairs 1 and 5; submetacentric pairs 2, 3, and 4; telocentric pairs 6, 7, 8, 9, 10, 11, and 12) (Fig. 1).

The karyotypes obtained in this study and those already published for the genus *Adenomera* are shown in Table 2 with their respective diploid number, fundamental number, and karyomorphology.

Discussion

The number of cytogenetic studies on anurans has grown in recent years (e.g., Ferro et al. 2018; Gazoni et al. 2018; Marciano-Jr et al. 2021); however, information for some families and/or genera is still scarce. The genus *Adenomera* comprises common and abundant species, some of which often occur syntopically (Cassini et al. 2020), but cytogenetic data for the entire genus are still scant compared to other anuran genera. So far, only six of the 29 described species have been karyotyped. Campos (2009) analyzed four populations of *Adenomera* from the state of São Paulo and identified two species, *Adenomera marmorata* and *Adenomera* aff. *bokermanni* with distinct karyotypes. *Adenomera marmorata* shows a variation in chromosome pair 12, which is




Figure 1. Karyotype of “*thomei*” clade specimens with conventional Giemsa staining **a** *Adenomera* cf. *thomei* from Ilhéus, Bahia, Brazil **b, c** *Adenomera* sp. L from Igrapiúna, Bahia, Brazil. All specimens showed the following karyomorphology: pairs 1 and 5 metacentric, 2–4 submetacentric, and 7–12 telocentric. Scale bar: 5 μ m.

metacentric in the populations of the state of São Paulo. Thus, Campos et al. (2009) hypothesized that it is an interpopulation variation, which was later confirmed by Cassini et al. (2020) in a taxonomic study on the group that integrated DNA sequences, morphology, and bioacoustics. The specimen identified by Campos et al. (2009) as *A. aff. bokermanni* was collected in the municipality of Santa Branca in the state of São Paulo, which is outside the current distribution of *A. bokermanni*, which is restricted to the southern region of the state of Paraná (Cassini et al. 2020).

The specimens analyzed in the present study were cytogenetically similar to those of *A. marmorata* and *A. aff. bokermanni* (Campos et al. 2009). Campos et al. (2009) found an unusual diploid number ($2n = 23$) when they described the karyotype of *A. aff. bokermanni*, Voucher - CFBH 11531, and concluded that it was most likely an indicative of a centric fusion involving the telocentric chromosome pairs 7 and 9. The authors

stated that it is not possible to determine with certainty whether the differences in chromosome pairs 7 and 9 correspond to a variation restricted to the specimen analyzed. Therefore, chromosome pairs 7 and 9 will not be used for comparison in our analyses. The chromosomes of pair 8 in the specimens analyzed in the present study are telocentric, whereas those of *A. aff. bokermanni* are subtelocentric (Campos et al. 2009).

Furthermore, the specimens of *Adenomera* cf. *thomei* (Ilhéus, BA) and *Adenomera* sp. L (Igrapiúna, BA) in the present study showed a karyotype ($2n = 24 - FN = 34$) identical to that of the specimen CFBH1512 from Santa Branca (SP) and the specimen CFBH 1713 (*Adenomera* sp. J). Moreover, no bioacoustic, molecular (DNA), or morphological data are available for the Ilhéus population and a taxonomic review including all species within the clade is needed to shed light on their specific limits.

Comparative cytogenetics can be considered an important tool for recovering phylogenetic relationships and confirming taxonomic identity (e.g., Baker 1970; Silva et al. 2004; Aguiar Jr et al. 2007; Urdampilleta et al. 2013; Cassini et al. 2020). The results presented here will contribute to expand the information on the taxonomy and phylogeny of the “*thomei*” clade and consequently lead to the delimitation of its taxa.

Acknowledgements

We thank Laisa Santos, Leildo Carilo and Fernanda Natascha for their help during field work. We also thank Laís Leal Lopes for her help in the laboratory and for kindly reviewing an earlier version of this manuscript. Thanks are also due to Carter Robert Miller for reviewing the manuscript. We also thank the kind input of two anonymous reviewers that improved the manuscript. We thank Reserva Ecológica Michelin for the support during field work. This study is part of the doctoral thesis by Ramon Costa Dominato at the Programa de Pós-Graduação em Genética e Biologia Molecular, Universidade Estadual de Santa Cruz, Ilhéus, Bahia, Brazil. We also thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the scholarship to Ramon Costa Dominato. Cléa dos Santos Ferreira Mariano, Janisete Gomes Silva, and Victor Goyannes Dill Orrico are fellows of the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

References

- Aguiar Jr O, Bacci Jr M, Lima AP, Rossa-Feres DDC, Haddad CFB, Recco-Pimentel SM (2007) Phylogenetic relationships of *Pseudis* and *Lysapsus* (Anura, Hylidae, Hylinae) inferred from mitochondrial and nuclear gene sequences. *Cladistics* 23(5): 455–463. <https://doi.org/10.1111/j.1096-0031.2007.00154.x>
- Almeida AP, Angulo A (2006) A new species of *Leptodactylus* (Anura: Leptodactylidae) from the state of Espírito Santo, Brazil, with remarks on the systematics of associated populations. *Zootaxa* 1334(1): 1–25. <https://doi.org/10.11646/zootaxa.1334.1.1>

- Angulo A (2004) The evolution of the acoustic communication system in members of the genus *Adenomera* (Anura: Leptodactylidae): A comparative approach. Doctoral thesis. University of Toronto, Toronto, 232 pp.
- Angulo A, Cocroft RB, Reichle S (2003) Species identity in the genus *Adenomera* (Anura: Leptodactylidae) in southeastern Peru. *Herpetologica* 59(4): 490–504. <https://doi.org/10.1655/20-104>
- Baker RJ (1970) Karyotypic trends in bats. In: Wimsatt WA (Ed.) *Biology of bats*. Academic Press Inc, New York, 65–97. <https://doi.org/10.1016/B978-0-12-758001-2.50007-1>
- Bogart JP (1970) Systematic problems in the amphibian family Leptodactylidae (Anura) as indicated by karyotypic analysis. *Cytogenetic and Genome Research* 9(5): 369–383. <https://doi.org/10.1159/000130106>
- Bogart JP (1974) A karyosystematic study of frogs in the genus *Leptodactylus* (Anura: Leptodactylidae). *Copeia* 3: 728–737. <https://doi.org/10.2307/1442686>
- Campos RC, Ananias JF, Aguirre BC, Yamamoto M, Haddad CFB, Kasahara S (2009) Chromosome evolution in three Brazilian *Leptodactylus* species (Anura, Leptodactylidae), with phylogenetic considerations. *Hereditas* 146(2): 104–111. <https://doi.org/10.1111/j.1601-5223.2009.02100.x>
- Carvalho TR, Moraes LJ, Lima AP, Fouquet A, Peloso PL, Pavan D, Drummond LO, Rodrigues MT, Giaretta AA, Gordo A, Neckel-Oliveira S, Haddad CF (2021) Systematics and historical biogeography of Neotropical foam-nesting frogs of the *Adenomera heyeri* clade (Leptodactylidae), with the description of six new Amazonian species. *Zoological Journal of the Linnean Society* 191(2): 395–433. <https://doi.org/10.1093/zoolinnean/zlaa051>
- Cassini CS, Taucce PP, Carvalho TR, Fouquet A, Solé M, Haddad CF, Garcia PC (2020) One step beyond a broad molecular phylogenetic analysis: Species delimitation of *Adenomera marmorata* Steindachner, 1867 (Anura: Leptodactylidae). *PloS ONE* 15(2): e0229324. <https://doi.org/10.1371/journal.pone.0229324>
- De Sá RO, Grant T, Camargo A, Heyer WR, Ponssa ML, Stanley E (2014) Systematics of the neotropical genus *Leptodactylus* Fitzinger, 1826 (Anura: Leptodactylidae): phylogeny, the relevance of non-molecular evidence, and species accounts. *South American Journal of Herpetology* 9(s1). <https://doi.org/10.2994/SAJH-D-13-00022.1>
- Duellman WE (2005) *Cusco Amazónico. The Lives of Amphibians and Reptiles in an Amazonian Rainforest*. Cornell University Press, Ithaca, [xv +] 433 pp.
- Ferro JM, Taffarel A, Cardozo D, Grosso J, Puig MP, Suárez P, Akmentins MS, Baldo D (2016) Cytogenetic characterization and B chromosome diversity in direct-developing frogs of the genus *Oreobates* (Brachycephaloidea, Craugastoridae). *Comparative Cytogenetics* 10(1): 141–156. <https://doi.org/10.3897/CompCytogen.v10i1.5718>
- Fouquet A, Cassini SC, Haddad FBC, Pech N, Trefaut RM (2014) Species delimitation, patterns of diversification and historical biogeography of the neotropical frog genus *Adenomera* (Anura, Leptodactylidae). *Journal of Biogeography* 41(5): 855–870. <https://doi.org/10.1111/jbi.12250>
- Gazoni T, Haddad CFB, Narimatsu H, Cabral-De-Mello DC, Lyra ML, Parise-Maltempi PP (2018) More sex chromosomes than autosomes in the Amazonian frog *Leptodactylus pentadactylus*. *Chromosoma* 127(2): 269–278. <https://doi.org/10.1007/s00412-018-0663-z>

- Green DM, Sessions SK (1991) Nomenclature for chromosomes. In: Green DM, Sessions SK (Eds) Amphibian cytogenetics and evolution. Academic Press, San Diego, 431–432. <https://doi.org/10.1016/B978-0-12-297880-7.50021-4>
- Kuramoto M (1990) A list of chromosome numbers of anuran amphibians. Bulletin of Fukuoka University of Education 39: 83–127.
- Lourenço LB, Bacci-Júnior M, Martins VG, Recco-Pimentel SM, Haddad CFB (2008) Molecular phylogeny and karyotype differentiation in *Paratelmatobius* and *Scythrophrys* (Anura, Leptodactylidae). Genetica 132(3): 255–266. <https://doi.org/10.1007/s10709-007-9169-y>
- Lourenço LB, Targueta CP, Baldo D, Nascimento J, Garcia PC, Andrade GV, Haddad CFB, Recco-Pimentel SM (2015) Phylogeny of frogs from the genus *Physalaemus* (Anura, Leptodactylidae) inferred from mitochondrial and nuclear gene sequences. Molecular Phylogenetics and Evolution 92: 204–216. <https://doi.org/10.1016/j.ympev.2015.06.011>
- Marciano-Jr E, Tanajura G, Costa MA, Orrico VG, Dias IR (2021) Advertisement call and cytogenetics of *Xenohyla eugenioi* Caramaschi, 1998 (Amphibia; Anura; Hylidae), an enigmatic frog from northeastern Brazil. South American Journal of Herpetology 20(1): 17–23. <https://doi.org/10.2994/SAJH-D-18-00035.1>
- Menin M, Waldez F, Lima AP (2008) Temporal variation in the abundance and number of species of frogs in 10,000 ha of a forest in Central Amazonia, Brazil. South American Journal of Herpetology 3(1): 68–81. [https://doi.org/10.2994/1808-9798\(2008\)3\[68:TVITAA\]2.0.CO;2](https://doi.org/10.2994/1808-9798(2008)3[68:TVITAA]2.0.CO;2)
- Pyron RA, Wiens JJ (2011) A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of extant frogs, salamanders, and caecilians. Molecular phylogenetics and evolution 61(2): 543–583. <https://doi.org/10.1016/j.ympev.2011.06.012>
- Silva APZ, Garcia P, Martins V, Bacci M, Kasahara S (2004) Chromosomal and molecular analyses of *Leptodactylus gracilis gracilis*, *L. gracilis delattini*, and *L. plaumanni* (Anura, Leptodactylidae): taxonomic implications. Amphibia-Reptilia 25(2): 185–196. <https://doi.org/10.1163/1568538041231193>
- Schmid M (1978) Chromosome banding in Amphibia. Chromosoma 66(4): 361–388. <https://doi.org/10.1007/BF00328536>
- Suárez P, Cardozo D, Baldo D, Pereyra MO, Faivovich J, Orrico VGD, Catroli GF, Grabiele M, Bernarde PS, Nagamachi CY, Haddad CFB, Pieczarka JC (2013) Chromosome evolution in Dendropsophini (Amphibia, Anura, Hylinae). Cytogenetic and Genome Research 141(4): 295–308. <https://doi.org/10.1159/000354997>
- Targueta CP, Rivera M, Souza MB, Recco-Pimentel SM, Lourenço LB (2010) Cytogenetic contributions for the study of the Amazonian *Engystomops* (Anura; Leiuperidae) assessed in the light of phylogenetic relationships. Molecular Phylogenetics and Evolution 54(3): 709–725. <https://doi.org/10.1016/j.ympev.2009.10.018>
- Urdampilleta JD, Coulleri JP, Ferrucci MS, Forni-Martins ER (2013) Karyotype evolution and phylogenetic analyses in the genus *Cardiospermum* L. (Paullinieae, Sapindaceae). Plant Biology 15(5): 868–881. <https://doi.org/10.1111/j.1438-8677.2012.00679.x>
- Zaracho VH, Hernando AB (2011) The karyotype of *Adenomera diptyx* (Boettger 1885) (Anura, Leptodactylidae) from northeastern Argentina. Genetics and Molecular Biology 34(1): 84–87. <https://doi.org/10.1590/S1415-47572010005000116>

ORCID

Ramon Costa Dominato <https://orcid.org/0000-0003-1939-1374>

Guilherme Costa de Oliveira <https://orcid.org/0000-0001-9990-9633>

Carla Santana Cassini <https://orcid.org/0000-0003-0366-645X>

Victor Goyannes Dill Orrico <https://orcid.org/0000-0002-4560-4006>

Cléa dos Santos Ferreira Mariano <https://orcid.org/0000-0001-7188-8201>

Janisete Gomes Silva <https://orcid.org/0000-0002-5570-5484>